



QUASIMEME

Quality assurance of information
for marine environmental monitoring

Certificate of Analysis



DSP shellfish toxins

REFERENCE MATERIAL

BT11 sample 9



Certificate of Analysis BT11 9

General Information

In this report an overview is given of analytical data for this sample collected in our proficiency testing program. The consensus values are calculated using a robust statistical model. With this NDA model mean and standard deviation are calculated using all reported data when at least 4 results are left after removal of reported 'lower than' (<) and 0 (= zero) values. No outliers are removed.

This report is divided into two sections: Consensus Values and Indicative Values. The division is made on the reliability of the data. Consensus Values are based on at least 10 results while the relative uncertainty is smaller than 6.25%. Indicative Values are based on a relative uncertainty of maximum 35% with at least 4 and less than 10 results or a relative uncertainty higher than 6.25%.

For each determinand the following parameters are given: mean, standard deviation, coefficient of variation, number of results, median, MAD (Median of Absolute Deviation) and the uncertainty in the assigned value. The confidence limits (at 95 % probability) are calculated for these determinands.

The results of each determinand is expressed on a wet weight basis.

Sample information

QUASIMEME reference materials cover a range of natural Shellfish toxins species from contaminated waters from the North Sea and/or Mediterranean.

This BT11 sample 9 of Mussel tissue (*Mytilus Edulis*) from Marine Institute, Ireland is prepared for the QUASIMEME proficiency programs. The results on which the values in this report are based were taken from the periods given in the following table.

Year.Round	Program	Sample Round Id
2016.1	BT11	QST208BT
2014.2	BT11	QST181BT



Consensus Values BT11

Method: Toxins(SF) - BT11

Element	Unit	Mean	Std.Dev.	CV %	N	Median	MAD	Uncertainty	95 % confidence limits		
AZA-1	µg/kg	133	23.1	17.4	68	135	15.9	3.5	127	-	139
AZA-2	µg/kg	44.8	8.11	18.1	60	45.0	5.57	1.31	42.7	-	46.9
AZA-3	µg/kg	33.7	7.24	21.5	57	34.0	4.90	1.20	31.8	-	35.7
AZA-total	µg AZA eq./kg	255	41.9	16.4	59	252	29.1	6.8	244	-	266
Free-DTX2	µg/kg	528	99.6	18.9	66	513	66.8	15.3	504	-	553
free-Okadaic-Acid	µg/kg	98.8	27.40	27.7	66	101.3	18.90	4.22	92.1	-	106
Total-free-OA+DTX1+DTX2	µg OA eq./kg	431	92.8	21.5	64	421	62.6	14.5	408	-	454
Total-DTX2	µg/kg	831	175.2	21.1	61	848	117.7	28.0	786	-	876
Total-Okadaic-Acid	µg/kg	238	42.6	17.9	62	236	29.0	6.8	227	-	248
Total-hy-OA+DTX1+DTX2	µg OA eq./kg	745	131.7	17.7	61	768	87.5	21.1	711	-	779
Total OA group + PTX group	µg OA eq./kg	737	148.3	20.1	46	734	99.3	27.3	693	-	781
YTX	mg/kg	1.20	0.175	14.6	59	1.20	0.120	0.029	1.16	-	1.25
Total-YTX	mg YTX eq./kg	1.20	0.164	13.7	49	1.18	0.110	0.029	1.15	-	1.25