

WAGENINGEN EVALUATING PROGRAMS FOR ANALYTICAL LABORATORIES

Certificate of Analysis



International Sediment Exchange for Tests on Organic Contaminants

REFERENCE MATERIAL
SETOC sample 704





Certificate of Analysis SETOC 704

General Information

In this report an overview is given of analytical data for this sample collected in our proficiency testing program. The consensus values are calculated using a robust statistical model. With this NDA model mean and standard deviation are calculated using all reported data when at least 8 results are left after removal of reported 'lower than' (<) and 0 (= zero) values. No outliers are removed.

This report is divided into three sections: Consensus Values, Indicative Values and Values for Information. The division is made on the reliability of the data. Consensus Values are based on at least 16 results while the coefficient of variation is smaller than 25 %. Indicative Values are based on at least 8 and less than 16 results or a coefficient of variation between 25 % and 50 %. Other values, based on more than 2 and less than 8 results or a coefficient of variation higher than 50 %, are given for information only.

In the sections with Consensus Values and Indicative Values the following parameters are given: mean, standard deviation, coefficient of variation, number of results, median and MAD (Median of Absolute Deviation) and the uncertainty in the consensus values. The confidence limits (at 95 % probability) are calculated for these determinands.

In the section with Information Values the following parameters are given: median, MAD and number of results. For determinands which have at least 5 results reported as smaller than (<) the median of these 'smaller than results' is calculated. In some cases this median of '<' values is much smaller than median and mean of the indicative values. This may be caused by a too optimistic (too low) value for the detection limit reported by a (small) majority of participating laboratories who report '<'-values.

All values, expressed on a weight basis (kg or %), are reported in oven dry (105 °C) material. Moisture is reported in the material as received.

Sample information

WEPAL reference materials are from natural sources only. There is no spiking, mixing or other alterations of the samples. For sample preparation the SETOC samples are dried at 40 $^{\circ}$ C and milled to pass a 0.5 mm sieve.

This SETOC sample 704 of Sediment from Netherlands is prepared for the WEPAL proficiency programs. The sample is used in 1 period (or round). The results on which the values in this report are based were taken from the period given in the following table.

Year	Round	Number
1992	4	1



Consensus Values SETOC 704



Method: Polycyclic aromatic hydrocarbons

Element benzo(b)fluoranthene Std.Dev. Uncertainty Unit CV % Ν Median MAD 95 % confidence limits Mean μg/kg 90.6 21.45 23.7 20 90.0 15.00 5.99 80.6 -101



Zn

Indicative Values SETOC 704



180

220

			alivo raid	· •	• •					
Method: Polycyclic aromatic hydro	carbons									
Element	Unit	Mean	Std.Dev.	CV %	N	Median	MAD	Uncertainty	95 % confiden	
anthracene	μg/kg	14.9	6.72	45.3	11	16.0	5.00	2.53	10.4 -	19.3
benzo(a)pyrene	μg/kg	52.9	26.08	49.3	18	48.0	16.00	7.68	40.0 -	65.8
benzo(ghi)perylene	μg/kg	58.2	23.34	40.1	19	60.0	15.50	6.69	47.0 -	69.4
benzo(k)fluoranthene	μg/kg	46.3	21.91	47.3	17	50.0	16.00	6.64	35.1 -	57.5
chrysene	μg/kg	51.7	17.86	34.6	19	52.0	12.00	5.12	43.1 -	60.3
fluoranthene	μg/kg	111	30.1	27.1	20	115	21.0	8.4	96.8 -	125
pyrene	μg/kg	76.3	31.23	40.9	16	78.0	22.00	9.76	59.8 -	92.9
Method: Polychlorobiphenyls										
Element	Unit	Mean	Std.Dev.	CV %	N	Median	MAD	Uncertainty	95 % confiden	ce limits
PCB 101	μg/kg	2.03	0.528	25.9	8	2.00	0.350	0.233	1.60 -	2.46
PCB 138	μg/kg	4.44	1.787	40.3	8	4.80	1.300	0.790	2.98 -	5.89
PCB 153	μg/kg	4.13	1.488	36.0	8	4.55	1.100	0.658	2.92 -	5.34
PCB 180	μg/kg	2.12	0.489	23.0	8	2.19	0.350	0.216	1.73 -	2.52
Method: Metals										
Element	Unit	Mean	Std.Dev.	CV %	N	Median	MAD	Uncertainty	95 % confiden	ce limits
As	mg/kg	10.7	1.44	13.5	10	10.8	1.00	0.57	9.71 -	11.7
Cd	mg/kg	0.859	0.1468	17.1	12	0.865	0.0950	0.0530	0.766 -	0.95
Cr	mg/kg	26.5	5.01	18.9	11	28.5	3.70	1.89	23.1 -	29.8
Cu	mg/kg	14.4	2.82	19.6	12	15.0	2.00	1.02	12.6 -	16.2
Hg	mg/kg	0.301	0.0258	8.6	9	0.310	0.0200	0.0108	0.282 -	0.32
Ni	mg/kg	15.6	4.35	27.9	11	15.0	3.20	1.64	12.7 -	18.4
Pb	mg/kg	36.8	8.83	24.0	12	36.4	6.10	3.19	31.2 -	42.3
	·									

15.9

12

203

20.9

11.5

200

31.9

mg/kg





Informative Values SETOC 704

Method: Polycyclic aron	natic hydrod		Results smaller than (<)			
Element	Unit	Median	MAD	N	Median of <	N
acenaphthene	μg/kg	17.8	4.25	6	50.0	13
acenaphthylene	μg/kg	-	-	0	30.0	16
benz(a)anthracene	μg/kg	46.5	19.00	16	100.0	6
dibenz(ah)anthracene	μg/kg	20.0	6.00	7	40.0	12
fluorene	μg/kg	30.4	15.38	16	100.0	5
indeno(1,2,3-cd)pyrene	μg/kg	70.0	30.00	19		
naphthalene	μg/kg	51.0	35.00	9	35.0	10
phenanthrene	μg/kg	95.0	43.00	20		

Method: Polychlorobiphenyls					Results smaller t	Results smaller than (<)		
Element	Unit	Median	MAD	N	Median of <	N		
PCB 028	μg/kg	2.75	0.750	6	5.00	10		
PCB 052	μg/kg	1.73	0.725	6	6.00	10		
PCB 118	µg/kg	3.90	0.100	4	8.00	10		

Method: Organochlori	Results smaller ti	han (<)				
Element	Unit	Median	MAD	N	Median of <	N
aldrin	μg/kg	-	-	0	10.0	13
beta-HCH	μg/kg	-	-	0	8.00	13
delta-HCH	μg/kg	-	-	0	5.50	6
dieldrin	μg/kg	-	-	0	10.0	13
endrin	μg/kg	-	-	0	4.50	14
gamma-HCH	μg/kg	-	-	0	5.00	13
hexachlorobenzene	μg/kg	1.40	0.400	5	10.00	10
o,p`-DDD	μg/kg	-	-	0	7.50	10
p,p`-DDD	μg/kg	2.00	0.290	3	10.00	10
p,p`-DDE	μg/kg	1.000	-	5	10.000	11
p,p`-DDT	μg/kg	1.60	0.600	3	10.00	12